



CrossMark
click for updates

Research

Cite this article: Rando JC, Pieper H, Alcover JA. 2014 Radiocarbon evidence for the presence of mice on Madeira Island (North Atlantic) one millennium ago. *Proc. R. Soc. B* **281**: 20133126.
<http://dx.doi.org/10.1098/rspb.2013.3126>

Received: 28 November 2013

Accepted: 14 January 2014

Subject Areas:

ecology, palaeontology

Keywords:

AMS ^{14}C , *Mus musculus*, island colonization, extinction, human presence, oceanic island

Author for correspondence:

Josep Antoni Alcover

e-mail: jaalcover@imedea.uib-csic.es

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2013.3126> or via <http://rspb.royalsocietypublishing.org>.

Radiocarbon evidence for the presence of mice on Madeira Island (North Atlantic) one millennium ago

Juan Carlos Rando¹, Harald Pieper² and Josep Antoni Alcover^{3,4}

¹Departamento de Biología Animal (Zoología), Universidad de La Laguna Tenerife, Canary Islands, Spain

²Ulmenstrasse 21, Schwentinal 24223, Germany

³Departament de Biodiversitat i Conservació, IMEDEA (CSIC-UIB), Cr Miquel Marqués 21, Esporles, Mallorca, Balearic Islands 07190, Spain

⁴Department of Mammalogy, American Museum of Natural History, New York, NY 10024-5192, USA

Owing to the catastrophic extinction events that occurred following the Holocene arrival of alien species, extant oceanic island biotas are a mixture of recently incorporated alien fauna and remnants of the original fauna. Knowledge of the Late Quaternary pristine island faunas and a reliable chronology of the earliest presence of alien species on each archipelago are critical in understanding the magnitude and tempo of Quaternary island extinctions. Until now, two successive waves of human arrivals have been identified in the North Atlantic Macaronesian archipelagos (Azores, Madeira, Selvagens, Canary and Cape Verde Islands): ‘aboriginal’, which is limited to the Canary Islands around two millennia ago, and ‘colonial’, from the fourteenth century onwards. New surveys in Ponta de São Lourenço (Madeira Island) have allowed us to obtain and date ancient bones of mice. The date obtained (1033 ± 28 BP) documents the earliest evidence for the presence of mice on the island. This date extends the time frame in which the most significant ecological changes occurred on the island. It also suggests that humans could have reached Madeira before 1036 cal AD, around four centuries before Portugal officially took possession of the island.

1. Introduction

Oceanic islands provide important insights for the study of evolution. Owing to the fact that catastrophic extinction events occurred everywhere following the Holocene arrival of alien species, extant island biotas are a mixture of the recently incorporated alien fauna and the remnants of the original fauna. Evolution and extinction have been intensively studied in island vertebrates [1–3]. The North Atlantic Macaronesian archipelagos (i.e. Azores, Madeira, Selvagens, Canary and Cape Verde Islands; figure 1), with different numbers of islands, geological ages, distances to the mainland and timing of human arrival, are ideal territories for the study of such processes. The knowledge of both Late Quaternary pristine island faunas and a reliable chronology of the early presence of aliens on each archipelago are critical to understanding the magnitude and tempo of Quaternary island extinctions.

Accelerator mass spectrometry (AMS) radiocarbon dating on commensal rodents has been widely used as a powerful tool to establish accurate chronologies for the early human presence on islands [4,5]. Until now, two successive waves of human colonization have been identified in Macaronesia: ‘aboriginal’ and ‘colonial’. In Macaronesia, only the Canary Islands harboured ‘aboriginal’ populations. AMS ^{14}C dating on introduced house mouse *Mus musculus domesticus* [6], and on endemic fauna of sites without it, indicates that humans reached this archipelago sometime between 756 cal BC and 313 cal AD [7]. On the other hand, the ‘colonial’ colonization wave affected all the Macaronesian archipelagos and took place from the fourteenth century onwards.

Current data indicate major extinction events in the Macaronesian islands subsequent to human arrivals. Among Upper Pleistocene–Holocene original




Figure 1. Location of the Macaronesian archipelagos. The fossil site where the bones of mice were collected is indicated on Madeira Island (red arrow). (Online version in colour.)

faunas of the Canary Islands, two genera and three species of endemic rodents, one species of giant lizard and at least seven species of endemic birds (two shearwaters, one quail, one oystercatcher, one bunting and two finches) and five species of non-endemic birds are now extinct. Chronological data indicate that all these species became extinct after human arrival [8–13]. A recent and impressive loss of endemic species seems to have occurred in the Madeira archipelago [14]. In this small archipelago, with only two main islands (Madeira and Porto Santo) and without terrestrial native mammals (excluding bats), bones obtained from Holocene dune fossil sites indicate that at least two-thirds of the endemic birds

(three species of rail, two quails, one scops owl, one thrush and one finch), plus two non-endemic species became extinct [14,15]. In electronic supplementary material, table S1, we present the list of the extinct birds and mammals from the Macaronesian archipelagos.

According to historical data, in the year 1419 the Portuguese took possession of Madeira, and the island was subsequently populated. It is usually assumed that, very probably, the Madeiran extinctions took place within the past 600 years. On Madeira, two different lines of evidence point to the presence of mice previous to the Portuguese colonization: (i) the finding of four subfossil bones of mice *Mus*




Figure 2. (a) Radiocarbon calibrated age range for the ancient mice from Madeira (upper and lower limits of 2σ confidence interval), Viking Age [22], proposed colonization time interval from mitochondrial D-loop sequences of current house mouse population of Madeira [23], and the date for Portuguese presence on the island. (b) Skull of ancient mouse (ventral view) in Ponta de São Lourenço (Madeira Island). (Online version in colour.)

musculus s.l. in a dune deposit [16], although the lack of radiocarbon dating on these materials did not allow us to confirm an age prior to the fifteenth century; (ii) the mtDNA haplotypes from current populations of house mouse *M. m. domesticus* of Madeira, which show similarities with those of Scandinavia and northern Germany (but not with the Portuguese mainland). This second line of evidence suggests that northern Europe was the source area, and raises the intriguing possibility that the Vikings could have brought the house mouse to the island [17], although it should be noted that to date there are no historical references of Viking voyages to Macaronesia.

The house mouse *M. musculus* spread from southeast Asia through Eurasia with human migrations during the Holocene. Two subspecies are currently present in Europe: the Western house mouse *M. m. domesticus*, in the Mediterranean and Western Europe; and the Eastern house mouse *M. m. musculus*, in Central and Eastern Europe [18]. The expansion of the Western house mouse in the Mediterranean began in the Eastern basin around the eighth millennium BC. It colonized the Western Mediterranean and northern Europe during the first millennium BC [19]. The Western house mouse was well settled on the Northeast coast of Spain in the second half of the first millennium BC [20]. The Eastern house mouse has been present in Eastern Europe at least since the fourth millennium BC [21].

The aim of this paper is to provide the first AMS ¹⁴C date of ancient mouse bones from Madeira, and to point out the possible consequences of this new date.

2. Material and methods

(a) New ancient samples of mouse bones

The new surveys on Madeira have provided two small samples of mouse bones from a dune site in Ponta de São Lourenço (red arrow in figure 1). In May 2010, a first sample (a single right mandible of 0.04 g) was collected. In November of 2011, at the same place, a second sample was collected. It included a fragmented skull (figure 2b), two fragmented jaws, three tibia, one complete femur, one fragmented femur, two fragmented humeri, two fragmented pelvises, three isolated incisors and

one caudal vertebra (total weight = 0.42 g). Other taxa present at this site are shearwaters (*Puffinus* sp.), petrels (*Pterodroma* sp.), pigeons (*Columba* sp.), an undescribed species of quail (*Coturnix*), the Madeiran scops owl (*Otus mauli*), the barn owl (*Tyto alba*) and several undescribed species of passerines (at least *Turdidae* and *Fringillidae*) [14,15].

(b) Radiometric dating and isotopic methods

Samples were dated through AMS at the Radiocarbon Accelerator Unit of the University of Oxford. AMS determinations were made on extracts of ultrafiltered bone gelatin [24]. Additionally, a fragment of barn owl humerus and three bones (a complete femur and tibiotarsus and a fragmented tarsometatarsus) of a new, still undescribed, species of quail from the same locality were AMS dated using an improved collagen extraction [25] by modified Longin method [26] at the Koninklijk Instituut voor Het Kunstpatrimonium (Brussels). No difficulties in obtaining the ¹⁴C date were reported from either laboratory.

Isotope ratio for C are presented as δ values, where $\delta = 1000[(R_{\text{sample}}/R_{\text{standard}}) - 1]$, and $R = {}^{13}\text{C}/{}^{12}\text{C}$. The isotopic reference standard is Vienna–PeeDee Belemnite.

(c) Conventions used for radiometric ages

The AMS radiocarbon dates given in years BP are conventional radiocarbon ages with standard error, where BP is before present (the year 1950), following standard reporting procedures. Radiocarbon calibration was calculated using the software OxCal v. 4.2 [27] and the IntCal13 dataset [28]. Dates calculated from calibration are expressed as 2σ intervals (95.4% confidence) and given as 'cal AD' or 'cal BC', AD being, *Annus Domini* and BC, *Before Christ*.

3. Results

Owing to their fragmentary condition and to the absence of any other rodent of Quaternary age in the island, we have cautiously attributed these new mouse bones to *M. musculus s.l.*

The first sample submitted for AMS-dating turned out to be too small to furnish a ¹⁴C age. The second sample (Lab-Code: OxA-26216) delivered the earliest testimony of the presence of mice on the island known so far: 1033 ± 28 BP, with a 2σ confidence interval of 903–1036 cal AD (figure 2a). A barn owl bone from the same site offers a similar

age (Lab-Code KIA-48920: 1060 ± 30 BP; 897–1024 cal AD), whereas a quail turned out to be older (Lab-Code KIA-47430: 2755 ± 55 BP; 1021–806 cal BC).

The $\delta^{13}\text{C}$ value for the dated mouse sample (-17.66%) falls within the range of the two bones of terrestrial birds from the same site (-15.53% for the barn owl and -18.6% for the quail), and within the range of dated rodent samples from the Canary Islands obtained at several sites located far inland from the shore (-16.99 to -20.12% for house mouse and -15.98 to -18.62% for lava mouse *Malpaisomys insularis* in the Canary Islands) [7]. Samples of animals with a marine diet (i.e. shearwaters) from the Canary Islands display higher $\delta^{13}\text{C}$ values (-11.51 to -13.61%) [12,13]. See list of $\delta^{13}\text{C}$ data of Holocene bird and mammal bones from the Macaronesian islands in the electronic supplementary material, table S2.

Our data indicate that it is highly unlikely that the mouse bones from Madeira that have been dated reflect a marine component in the diet. It is consequently improbable that the calibrated age of this sample was affected by the ^{14}C marine reservoir effect.

4. Conclusion

Although a natural colonization by mice—via natural rafts, like those suggested for the presence of endemic rodents on the Canary Islands, which are located around 100 km from the African coast [29]—cannot be totally excluded, it appears to be highly improbable. It should be mentioned here that the Western Mediterranean islands, despite being closer to the mainland than Madeira, were not colonized by mice until the commercial and demographic spread of Phoenicians and Greeks during the last millennium BC [30]. Also, because of the island's isolation, it is highly unlikely that the ancient mouse sample reached Madeira transported by a bird (i.e. inside a pellet).

If we assume that humans transported mice to Madeira, the new date presented herein suggests that humans could have reached the island before 1036 cal AD, thus around four centuries before Portugal officially took possession of

the island. The 2σ confidence interval of OxA-26216 (900–1036 cal AD) falls within Viking Age (eighth to eleventh centuries) [22], and agrees with one of the two proposed colonization time intervals obtained from mitochondrial D-loop sequences of current house mouse populations of Madeira (the mismatch distribution of pairwise genetic differences approach: 910–1185 AD) [23]. mtDNA haplotypes from current mouse populations of Madeira show similarities with those of Scandinavia and northern Germany [17], and it has been reported that Vikings transported house mice to the places they reached [31,32]. All these data suggest, but do not prove, a relationship between the Viking voyages and the presence of *Mus* on Madeira.

Once settled, the mouse population could have reached very high densities owing to its high reproductive potential, particularly in the absence of rats (*Rattus* spp.), which limit their number. In this case, a wide range of native biota could have been impacted, and the magnitude of the impact could have been huge, as occurred on islands without rats [33,34]. Nevertheless, more radiocarbon, genetic and morphometric data [6,19–21] are necessary to establish the precise evolutionary history of mice on Madeira (including the subspecific diagnosis of the ancient mice), the precise extent of the impact of the arrival of mouse on the island, and the causes and precise chronologies of bird extinctions on Madeira. The new date provided herein demonstrates the early presence of mice on Madeira and provides a longer time span in which the most severe ecological changes during the Late Holocene occurred on the island after the arrival of mice (i.e. at least *ca* one millennium ago).

Acknowledgements. We thank Thomas Highman (Oxford), Mark Van Strydonck and Mathieu Boudin (Brussels) for their comments and information on dating procedures. We thank Hannah Bonner, Juan Carlos Illera, Paulo Oliveira, Dilia Meneses and Manuel Biscoito for their assistance with several aspects of this paper. The authors declare they have no competing interests.

Funding statement. This work was partially supported by DGICYT Research Project CGL2012-38089.

References

- Olson SL, James HF. 1982 Fossil birds from the Hawaiian Islands: evidence for wholesale extinction by man before Western contact. *Science* **217**, 633–635. (doi:10.1126/science.217.4560.633)
- Steadman DW. 1995 Prehistoric extinctions of pacific island birds: biodiversity meets zooarchaeology. *Science* **267**, 1123–1131. (doi:10.1126/science.267.5201.1123)
- Lossos JB, Ricklefs RE. 2009 Adaptation and diversification on islands. *Nature* **457**, 830–836. (doi:10.1038/nature07893)
- Wilmshurst JM, Anderson AJ, Higham TFG, Worthy TH. 2008 Dating the late prehistoric dispersal of Polynesians to New Zealand using the commensal Pacific rat. *Proc. Natl Acad. Sci. USA* **105**, 7676–7680. (doi:10.1073/pnas.0801507105)
- Wilmshurst JM, Hunt TL, Lipo CP, Anderson AJ. 2011 High-precision radiocarbon dating shows recent and rapid initial human colonization of East Polynesia. *Proc. Natl Acad. Sci. USA* **108**, 1815–1820. (doi:10.1073/pnas.1015876108)
- Michaux J, Cucchi T, Renaud S, García-Talavera F, Hutterer R. 2007 Evolution of an invasive rodent on an archipelago as revealed by molar shape analysis: the house mouse in the Canary Islands. *J. Biogeogr.* **34**, 1412–1425. (doi:10.1111/j.1365-2699.2007.01701.x)
- Alcover JA, Rando JC, García-Talavera F, Hutterer R, Michaux J, Trias M, Navarro JF. 2009 A reappraisal of the stratigraphy of Cueva del Llano (Fuerteventura) and the chronology of the house mouse (*Mus musculus*) introduction into the Canary Islands. *Palaeogeogr. Palaeocli. Palaeoecol.* **277**, 184–190. (doi:10.1016/j.palaeo.2009.03.016)
- Rando JC. 2003 Protagonistas de una catástrofe silenciosa: los vertebrados extintos de Canarias. *El Indiferente* **14**, 4–15.
- Rando JC, Alcover JA, Michaux J, Hutterer R, Navarro JF. 2012 Late-Holocene asynchronous extinction of endemic mammals on the Eastern Canary Islands. *Holocene* **22**, 801–808. (doi:10.1177/0959683611430414)
- Illera JC, Rando JC, Richardson DS, Emerson BC. 2012 Age, origin and extinctions of the avifauna of Macaronesia: a synthesis of phylogenetic and fossil information. *Quat. Sci. Rev.* **50**, 14–22. (doi:10.1016/j.quascirev.2012.07.013)
- Rando JC, Alcover JA, Illera JC. 2010 Disentangling ancient interactions: a new extinct passerine provides insights on character displacement among extinct and extant island finches. *PLoS ONE* **5**, e12956. (doi:10.1371/journal.pone.0012956)
- Rando JC, Alcover JA. 2008 Evidence for a second Western Palaearctic seabird extinction during the last Millennium: the Lava Shearwater *Puffinus*

- olsoni*. *Ibis* **150**, 188–192. (doi:10.1111/j.1474-919X.2007.00741.x)
13. Rando JC, Alcover JA. 2010 On the extinction of dune shearwater (*Puffinus holeae*) from the Canary Islands. *J. Ornithol.* **151**, 365–369. (doi:10.1007/s10336-009-0463-6)
 14. Pieper H. 1985 The fossil land birds of Madeira and Porto Santo. *Bocagiana* **88**, 1–6.
 15. Rando JC, Pieper H, Alcover JA, Olson SL. 2012 A new species of extinct fossil scops owl (Aves: Strigiformes: Strigidae: *Otus*) from the Archipelago of Madeira (North Atlantic Ocean). *Zootaxa* **3182**, 29–42.
 16. Pieper H. 1981 Ein subfossiles Vorkommen der Hausmaus (*Mus musculus s.l.*) auf Madeira. *Bocagiana* **59**, 1–3.
 17. Gündüz I, Auffray JC, Britton-Davidian J, Catalan J, Ganem G, Ramalhinho MG, Mathias ML, Searle JB. 2001 Molecular studies on the colonization of the Madeiran archipelago by house mice. *Mol. Ecol.* **10**, 2023–2029. (doi:10.1046/j.0962-1083.2001.01346.x)
 18. Auffray J-C, Marshall JT, Bonhomme F, Thaler L. 1990 Focus on the nomenclature of European species of *Mus*. *Mouse Genome* **88**, 7–8.
 19. Cucchi T, Vigne JD, Auffray JC. 2005 First occurrence of the house mouse (*Mus musculus domesticus* Schwarz & Schwarz, 1943) in the Western Mediterranean: a zooarchaeological revision of subfossil occurrences. *Biol. J. Linn. Soc.* **84**, 429–445. (doi:10.1111/j.1095-8312.2005.00445.x)
 20. Valenzuela-Lamas S, Baylac M, Cucchi T, Vigne J-D. 2011 House mouse dispersal in Iron Age Spain: a geometric morphometrics appraisal. *Biol. J. Linn. Soc.* **102**, 483–497. (doi:10.1111/j.1095-8312.2010.01603.x)
 21. Cucchi T, Balasescu A, Bem C, Radu V, Vigne J-D, Tresselt A. 2011 New insights into the invasive process of the Eastern house mouse (*Mus musculus musculus*): evidence from the burnt houses of Chalcolithic Romania. *Holocene* **21**, 1195–1202. (doi:10.1177/0959683611405233)
 22. White P. 2010 *Exploration in the world of the Middle ages, 500–1500*. New York, NY: Chelsea House Publishers.
 23. Förster DW, Gündüz I, Nunes AC, Gabriel S, Ramalhinho MG, Mathias ML, Britton-Davidian J, Searle JB. 2009 Molecular insights into the colonization and chromosomal diversification of Madeiran House Mice. *Mol. Ecol.* **18**, 4477–4494. (doi:10.1111/j.1365-294X.2009.04344.x)
 24. Bronk Ramsey C, Higham T, Bowles A, Hedges R. 2004 Improvements to the pretreatment of bone at Oxford. *Radiocarbon* **46**, 155–163.
 25. van Strydonck M, Boudin M, Ervynck A. 2005 Humans and *Myotragus*: the issue of sample integrity in radiocarbon dating. *Monogr. Soc. Hist. Nat. Balears.* **12**, 369–376.
 26. Brown TA, Vogel JS, Southon JR. 1998 Improved collagen extraction by modified Longin method. *Radiocarbon* **30**, 171–177.
 27. Bronk Ramsey C. 2009 Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**, 337–360.
 28. Reimer PJ *et al.* 2013 IntCal13 and Marine13 radiocarbon age calibration curves, 0–50,000 years cal BP. *Radiocarbon* **55**, 1869–1887. (doi:10.2458/azu_js_rc.55.16947)
 29. Pagès M, Chevret P, Gros-Balthazard M, Hughes S, Alcover JA, Hutterer R, Rando JC, Michaux J, Hänni C. 2012 Paleogenetic analyses reveal unsuspected phylogenetic affinities between mice and the extinct *Malpaisomys insularis*, an endemic rodent of the Canaries. *PLoS ONE* **7**, e31123. (doi:10.1371/journal.pone.0031123)
 30. Cucchi T, Vigne JD. 2006 Origin and diffusion of the house mouse in the Mediterranean. *Human Evol.* **21**, 95–106. (doi:10.1007/s11598-006-9011-z)
 31. Jones EP, Van Der Kooij J, Solheim R, Searle JB. 2010 Norwegian house mice (*Mus musculus musculus/domesticus*): distributions, routes of colonization and patterns of hybridization. *Mol. Ecol.* **19**, 5252–5264. (doi:10.1111/j.1365-294X.2010.04874.x)
 32. Jones EP, Skirnisson K, McGovern TH, Gilbert MTP, Willerslev E, Searle JB. 2012 Fellow travellers: a concordance of colonization patterns between mice and men in the north Atlantic region. *BMC Evol. Biol.* **12**, 35. (doi:10.1186/1471-2148-12-35)
 33. Angel A, Wanless RM, Cooper J. 2009 Review of impacts of the introduced house mouse on islands in the Southern Ocean: are mice equivalent to rats? *Biol. Invasions.* **11**, 1743–1754. (doi:10.1007/s10530-008-9401-4)
 34. Wanless RM, Ratcliffe N, Angel A, Bowie BC, Cita K, Hilton GM, Kritzing P, Ryan PG, Slabber M. 2012 Predation of Atlantic petrel chicks by house mice on Gough Island. *Anim. Conserv.* **15**, 472–479. (doi:10.1111/j.1469-1795.2012.00534.x)